

表2. Disk agar diffusion susceptibility 判定標準表

藥物名稱	濃度	供應商	判定標準(抑制圈大小-mm)		
			Resistant	Intermediate	Susceptible
Ampicillin	10µg	OXOID	≤ 13	14-16	≥ 17
Amoxicillin	20µg	OXOID	≤ 13	14-17	≥ 18
Cephalothin	30µg	OXOID	≤ 14	15-17	≥ 18
Ceftiofur	30µg	OXOID	≤ 17	18-20	≥ 21
Colistin	10µg	OXOID	≤ 8	9-10	≥ 11
Doxycycline	30µg	OXOID	≤ 12	13-15	≥ 16
Enrofloxacin	5µg	OXOID	≤ 16	17-22	≥ 23
Erythromycin	15µg	OXOID	≤ 13	14-22	≥ 23
Flofenicol	30µg	OXOID	≤ 14	15-18	≥ 19
Flumequin	30µg	OXOID	≤ 15	16-20	≥ 21
Gentamicin	10µg	OXOID	≤ 12	13-14	≥ 15
Kanamycin	30µg	OXOID	≤ 13	14-17	≥ 18
Neomycin	30µg	OXOID	≤ 12	13-16	≥ 17
Nalidixic acid	30µg	OXOID	≤ 13	14-18	≥ 19
Oxytetracycline	30µg	OXOID	≤ 15	16-20	≥ 21
Streptomycin	10µg	OXOID	≤ 11	12-14	≥ 15
Tetracycline	30µg	OXOID	≤ 14	15-18	≥ 19
Tiamulin	30µg	NEO-SENSITABS	≤ 7	8-9	≥ 10
Trimethoprim-Sulphamethoxazole	1.25/23.75 µg	OXOID	≤ 10	11-15	≥ 16
Tylosin	150µg	NEO-SENSITABS	≤ 22	23-25	≥ 26

(NCCLS, M31-A2)

表3. 藥物敏感性試驗對照表

建議選用(V)	名 稱	豬	建議選用(V)	名 稱	豬
	Ampicillin	R		Kanamycin	R
	Amoxicillin	R		Nalidixic acid	R
V	Colistin	S		Neomycin	R
V	Cephazolin	S		Oxytetracycline	R
	Doxycycline	R		Streptomycin	R
V	Enrofloxacin	S		Sulphamethoxazole/Trimethoprim	R
	Erythromycin	R		Tetracycline	R
	Florocol	R		Flumequine	R
	Gentamicin	R	V	Ceftiofur	S
	Tiamulin	R		Tylosin	R